


Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Children and Youth Services Review

journal homepage: www.elsevier.com/locate/chilyouth

Dosage matters: The relationship between participation in the Nurturing Parenting Program for infants, toddlers, and preschoolers and subsequent child maltreatment

Erin J. Maher^{a,*}, Lyscha A. Marcynyszyn^a, Tyler W. Corwin^b, Rhenda Hodnett^c

^a Casey Family Programs, 2001 Eighth Avenue, Suite 2700, Seattle, WA, 98121, USA

^b Northwest Social Research Group, 700 Broadway East, Unit 119, Seattle, WA, 98102, USA

^c Louisiana Office of Community Services, Department of Social Services, 627 N. Fourth Street 3–313, Baton Rouge, LA, 70802, USA

ARTICLE INFO

Available online 16 April 2011

Keywords:

Parent education
Maltreatment
Child welfare
Program evaluation

ABSTRACT

This article uses statewide data on caregivers of young children referred to the Nurturing Parenting Program (NPP) for allegations of abuse and neglect to examine the relationship between program dosage and subsequent maltreatment. At six months after participating in the program, caregivers who attended more sessions were significantly less likely to be reported for child maltreatment, holding other factors constant. At two years after participating, caregivers attending more sessions were significantly less likely to have a substantiated maltreatment incidence, controlling for other characteristics of families associated with maltreatment. These findings demonstrate the program is associated with preventing short-term allegations and longer-term substantiated incidences of maltreatment for a child welfare population. By demonstrating the importance of participation in the NPP, we increase the evidence about effective parent education programs for this population.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

The deleterious and long-term effects of maltreatment are well documented (Gilbert et al., 2009; Haapasalo & Pokela, 1999; Widom & Maxfield, 2001). Parent education, also called parent training, is one of the most commonly used interventions in child welfare to prevent child maltreatment (Barth et al., 2005; Huebner, 2002; Hurlburt, Barth, Leslie, & Landsverk, 2007). While the knowledge gap about these programs is shrinking (Waldfoegel, 2009), we know surprisingly little about the effectiveness of parent education for families involved in the child welfare system and about the implementation of these programs within the real-world constraints of child welfare agencies (Barth et al., 2005; Chaffin & Friedrich, 2004). These gaps are likely due to limited use of evidence-based parent education programs for families already involved in the child welfare system (Lee, Aos, & Miller, 2008) and limited monitoring of program implementation and outcomes, including longer-term follow-ups.

Casey Family Programs partnered with Louisiana's Department of Social Services to evaluate the effectiveness of one particular parent education program – the Nurturing Parenting Program (NPP) for Infants, Toddlers, and Preschoolers (Bavolek, 2005) – in preventing maltreatment. Program implementation began in late 2005 and in 2006 was fully implemented statewide. This study contributes to building an

evidence base of parent education in child welfare by evaluating a promising program model with a population of participants involved in the child welfare system following an allegation of abuse or neglect of one or more children in their care. In addition, it uses administrative data to document the changes in reported and substantiated incidences of post-intervention maltreatment over an extended period of time. In this study, we define “substantiated” as findings that abuse or neglect occurred as the result of a child welfare investigation. “Reported” refers to all investigated reports, which may or may not be substantiated.

1.1. Parent education programs

In child welfare, parent education serves families who are trying to keep their children from entering out-of-home care or who are seeking reunification (Barth et al., 2005). Yet, the use of evidence-informed or evidence-based parent education programs is scarce. For example, the parenting programs with the strongest evidence of overall effectiveness have most commonly been studied in clinical settings and primarily focused on child behavior problems (Barth et al., 2005). Parent training programs such as MultiSystemic Therapy (Henggeler, Schoenwald, Borduin, Rowland, & Cunningham, 1998), Parent-Child Interaction Therapy (PCIT) (Eyberg & Robinson, 1982), The Incredible Years (Webster-Stratton, 2007), and Parent Management Training (Patterson, Reid, & Eddy, 2002) are primarily focused on preventing, reducing, and treating children with serious behavior problems. These programs have been touted as having the most promise for use in child welfare based on their empirical evidence with other at-risk populations (Barth et al., 2005).

* Corresponding author. Tel.: +1 206 378 3317; fax: +1 866 519 3884.

E-mail addresses: emaher@casey.org (E. J. Maher), LMarcynyszyn@casey.org (L. A. Marcynyszyn), tyler.w.corwin@gmail.com (T. W. Corwin), Rhenda.Hodnett@la.gov (R. Hodnett).

While children with social and emotional difficulties represent a portion of the child welfare population and these programs are a valuable resource to meet their specific needs, just over 80% of all child victims of maltreatment are victims of neglect (including medical neglect), and one-third are under four years old (U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth & Families, Children's Bureau, 2010). Thus, questions of whether parent education can prevent maltreatment, particularly neglect, among young children, and, concomitantly, repeat maltreatment are paramount. And, effective interventions for parents with young children are especially critical as this age group is at greatest risk for entry into care (Wulczyn, Hislop, & Jones Harden, 2002). This period also sets the stage for positive parenting practices throughout a child's development and life course.

1.2. Programs for families at risk for entry into child welfare

Few randomized control trials of parent education programs in 'at risk' populations exist, and even fewer focus solely on families already involved in the child welfare system (for reviews see Barth et al., 2005; Johnson et al., 2008; Lundahl, Nimer, & Parsons, 2006). For families not involved in the child welfare system, a handful of prevention-focused, parenting education programs have been conducted using rigorous evaluation designs. Findings from these evaluations have been mixed. Many demonstrate positive effects on maltreatment outcomes (Britner & Reppucci, 1997; Bugental et al., 2002; Duggan et al., 2004; Huxley & Warner, 1993; Olds et al., 1997; Prinz, Sanders, Shapiro, Whitaker, & Lutzker, 2009), while others show inconclusive (DuMont et al., 2008; Wagner & Clayton, 1999) or no effects (Barth, 1991; Duggan et al., 1999; 2007; Marcenko, Spence, & Samost, 1996).

1.3. Programs for families in child welfare

To our knowledge, Chaffin et al. (2004) conducted the only published evaluation of parent training using a randomized control design with a child welfare population. They examined the effectiveness of PCIT in preventing maltreatment recurrence with 110 families. Their results indicated that PCIT reduced subsequent maltreatment among physically abusive parents more than two years later at follow-up. Despite these findings, child neglect outcomes were not improved by PCIT. A quasi-experimental design to evaluate Project SafeCare, which is an in-home parent training for families already involved in the child welfare system, has demonstrated promise for reducing maltreatment (Gershater-Molko, Lutzker, & Wesch, 2002), while other home-visiting programs with this population have not (MacMillan et al., 2005). Finally, several recent smaller-scale evaluations have studied the Incredible Years Parent Training Program with a child welfare population and examined intermediate variables hypothesized to be associated with child maltreatment, such as parenting attitudes (Letarte, Normandeau, & Allard, 2010; Marcynyszyn, Maher, & Corwin, 2011; Stern, Alaggia, Watson, & Morton, 2008), but there is a dearth of large-scale evaluations on maltreatment outcomes for this population. For a comprehensive review on this topic see studies by Lee et al. (2008); MacMillan et al. (2005); and Reynolds, Mathieson, and Topitzes (2009).

1.4. The Nurturing Parenting Program (NPP)

The NPP (Bavolek, 2005) is a parent education program identified as holding promise for effectiveness in child welfare. Based on the available research evidence and primary focus on reducing abusive or neglectful behavior, the California Evidence-Based Clearinghouse (CEBC) for Child Welfare rated the NPP as a Level 1 (highest level) for relevance to child welfare. However, despite findings from pre/post-test design evaluation studies (see Cowen, 2001; Devall, 2004; Hodnett, Faulk, Dellinger, & Maher, 2009), the lack of randomized control trials

resulted in a scientific rating of Level 3, "Promising Research Evidence," for evidence of effectiveness (CEBC for Child Welfare, 2010). Recently, the Substance Abuse and Mental Health Services Administration's National Registry of Evidence-Based Programs and Practices (SAMHSA NREPP) listed the NPP as a 3 out of 4 for quality of research and 4 out of 4 for readiness for dissemination (SAMHSA NREPP, 2010). In addition, the NPP contains all three of the essential components identified in a recent meta-analysis by Kaminski, Valle, Filene, and Boyle (2008) on parent training program effectiveness in other populations (e.g., not necessarily child welfare). These included teaching parents emotional communication and positive parent-child interaction skills, teaching about discipline consistency and time out, and providing parents with the opportunity to practice new skills with children during the training sessions (Kaminski et al., 2008).

1.5. Program dosage and recurrence of maltreatment

Parent participation and engagement in a program is pivotal for its effectiveness. Holding case characteristics believed to predict maltreatment recurrence constant (child vulnerability, family stress, partner abuse, and social support deficiencies), DePanfilis and Zuravin (2002) found that only attendance at services recommended in the case plans predicted less recurrence of child maltreatment. Further, a more recent review by Waldfoegel (2009) highlights the essential role of program dosage and family engagement as fundamental to the effectiveness of prevention programs such as parent education. As is especially true for a child welfare population, interventions must be of sufficient intensity and duration to address the severity of familial and sociodemographic risk factors (Lundahl et al., 2006; Thomlison, 2005). However, these findings conflict with earlier research by Littell (1997), which indicated that the length of family preservation services was unrelated to the recurrence of child maltreatment. These seemingly contradictory findings may be explained by recognizing an inherent confound. Those families considered higher risk are more likely to receive services over a longer period of time, but also are more likely to have additional maltreatment instances. At the same time, evidence also supports that higher risk families might be more difficult to engage in services for longer periods of time (Erickson & Egeland, 2011). Building on this research about program participation and engagement, the current evaluation examines whether dosage, as defined by the total number of NPP group or in-home sessions, is associated with maltreatment following program participation. We control for numerous factors known to be correlated with maltreatment, including whether a parent is at higher risk for repeat maltreatment (i.e., prior maltreatment history and parenting attitudes), which the literature supports is important to control for when examining the relationship between length of service, or, in our case, attendance and outcomes.

1.6. The current study

Although many studies identify maltreatment prevention as a program goal, most do not measure its occurrence. Instead, investigators tend to measure behaviors believed to be associated with neglect such as maladaptive parenting skills, social isolation, depression, or substance abuse. This study specifically examines the relationship between NPP dosage and maltreatment following program participation. Given that prior evaluation studies with families involved in the child welfare system have generally been low in methodological rigor due to selective sampling, small sample sizes, short follow-up periods, and lack of actual maltreatment outcome data (see Reynolds et al., 2009, for a review), the current evaluation is poised to make a much needed contribution to the field by examining maltreatment following NPP participation at both six months and two years post-intervention in a large, statewide child welfare population using state record data.

While we acknowledge that examining the relationship between NPP participation and post-intervention maltreatment using a cross-sectional design with follow-up and administrative data is not the gold standard in establishing effectiveness, it represents a step forward along a continuum to establish this program's promise for a child welfare population. Whereas this study does not fulfill the need for a randomized control trial, it builds on the existing knowledge base by analyzing the systematic, statewide implementation of one specific parent education program and provides an analysis of program dosage while holding factors associated with risk, such as caregiver education, age, marital status, and number of children in the household constant. Moreover, the current evaluation is conducted by investigators who are independent of the intervention developer.

Using administrative data merged with pre-intervention, caregiver self-report surveys on parenting behaviors and beliefs, this evaluation addresses the following research question: What is the relationship between NPP dosage and reported and substantiated incidences of child maltreatment following the program at six months and two years post-participation?

2. Method

2.1. Study population

Our study relies on the population of caregivers who attended the NPP in ten of the state's twelve family resource centers that served Louisiana's child welfare population between October 2005 and April 2008. Eleven centers offered the program, but families served in one of these eleven resource centers were excluded from the study due to the program modifications necessitated by Hurricane Katrina. With the exception of caregivers served at this one resource center, we used the total population of all caregivers served by the NPP during this time period — not a selected sample. The resource centers offered the NPP to all families with children under six with child abuse and/or neglect allegations, while screening for circumstances and behaviors that would prevent constructive participation in the program (serious cognitive impairment, work barriers, substance abuse), although these factors did not necessarily preclude participation. With the exception of a very small number of families who were screened out across the state (~50), all child welfare-involved families with infants, toddlers, or preschool-aged children in the state assessed as needing parent education were referred to the NPP. The study population consists of 564 caregivers who enrolled in the NPP at the ten resource centers. Eighteen caregivers participated in the NPP twice. Because the range of possible attendance differs for individuals who enrolled in the program twice, rather than only once, these caregivers were removed from the analysis, which resulted in 528 unique participants in the NPP.

2.2. The intervention

The NPP (Bavolek, 2005) is based on social learning theory (Bandura, 1977, 1986) and the associated premise that most parenting patterns are learned during childhood and replicated later in life when the child becomes a parent. The program is designed to assess, prevent, and treat maltreatment by developing nurturing parenting skills as a counter to the key constructs of abusive and neglectful parenting identified by Bavolek, Kline, McLaughlin, and Publicover (1979) from the literature and expert advisors (as cited in Bavolek, 2005). The constructs center around parental expectations of the child, empathy toward children's needs, use of corporal punishment as a means of discipline, parent-child role responsibilities, and children's power and independence.

The NPP is built on the core principle that empathy is the foundation of responsive parenting, for which there is general agreement that promoting nurturing and empathic parenting practices is critical to the safety and well-being of children (Donald & Jureidini, 2004; Kochanska

& Aksan, 1995; Laible, 2004). The specific program for parents of infants, toddlers, and pre school children focuses on parental self-awareness and empowerment, the development of empathy, understanding child development and the role of discipline, emotional communication, behavior skills training, the importance of nurturing routines, and making good choices for child safety (Bavolek & Dellinger-Bavolek, 1985). The program is designed to be flexible in its application and involves lesson guides, DVDs, parent handbooks, assessment inventories, behavioral modeling, discussion, role playing, and family activities to promote cognitive and affective learning. Finally, the NPP is structured to involve children in the learning process.

Louisiana's Department of Social Services provided funding to train resource center staff to become NPP facilitators. In addition, front-line child welfare staff were also trained so they could become familiar with the program model. The NPP was administered as a 16-week group-based program. Sixteen is the minimum number of sessions recommended for a child welfare population. Home visits were used as make-ups for missed sessions in order to maximize participation, as compliments to the group sessions to reinforce concepts for parents who needed more time, and as supplemental sessions for parents who had other needs. Group sessions lasted approximately two and a half hours, and home sessions averaged one hour. The NPP manual is written at a 5th grade level, and the state child welfare agency worked with the program developer to create an "Easy Reader" version of the materials for use with caregivers with more limited reading ability. To promote accessibility, the resource center staff were expected to assist caregivers in developing transportation plans as needed.

2.3. Data

Data for this analysis is drawn from three sources merged on unique case IDs: NPP attendance data; the state child welfare Tracking and Information Payment System (TIPS), which is the state's administrative child welfare data system; and, the Adult-Adolescent Parenting Inventory-2 (AAPI-2) pre- and post-test surveys (Bavolek & Keene, 2001). Attendance was recorded by the resource centers for each caregiver's participation in home and/or group sessions of the NPP. In addition to demographic information, TIPS data included up-to-date information on the dates, types, and status of child maltreatment investigations. The AAPI-2 pre-test and post-test surveys contain demographic questions and 40 statements, each measured on a 5-point Likert scale, to assess caregivers': (a) inappropriate expectations of children, (b) parental lack of empathy toward children's needs, (c) strong belief in the use of corporal punishment, (d) parent-child role reversal, and (e) oppression of children's power and independence.

2.4. Dependent variables

We examined the relationship between NPP participation and both reported and substantiated maltreatment. Federal child care statistics generally rely on counts of substantiated incidences. However, significant debate about the meaning and value of unsubstantiated reports exists in both the child welfare literature and among community stakeholders, and some research has found no difference in child well-being outcomes between families with substantiated and unsubstantiated reports (Hussey et al., 2005) or in risk of recidivism (Drake, Jonson-Reid, Way, & Chung, 2003). We also know that some unsubstantiated reports do, in fact, involve maltreatment (Waldfoegel, 2009), and many of the service needs among families with substantiated and unsubstantiated reports are similar. Nonetheless, given this debate and the fact that both substantiated and unsubstantiated reports can warrant interventions from and involve significant costs to child welfare, we chose to examine all reports and substantiated reports separately to allow for the possibility of differential associations with program participation.

The dependent variables for the four analytical models were whether or not a caregiver had a reported or substantiated incidence of maltreatment six months and two years subsequent to the end of their participation in the NPP. These variables were constructed from the TIPS data and included any allegation of neglect, physical abuse, sexual abuse, or other type of maltreatment during the relevant time period. The maltreatment allegation after participation in the NPP could be for any child under the caregiver's supervision, not necessarily for the child under six years, for whom they were referred to the program. The four dichotomous dependent variables were: (a) a reported incidence of child maltreatment within six months of the NPP closure; (b) a substantiated incidence of child maltreatment within six months of the NPP closure; (c) a reported incidence of child maltreatment within two years of the NPP closure; and (d) a substantiated incidence of child maltreatment within two years of the NPP closure.

The six-month window of opportunity for maltreatment after participation in the NPP was selected based upon the federal definition of maltreatment recurrence (Data Measures, Data Composites, National Standards To Be Used in the Child and Family Services Reviews, 2006). Fluke, Shusterman, Hollinshead, and Yuan (2005) demonstrate that while rates of repeat and reported maltreatment taper after six months following an incidence of child abuse and/or neglect, the cumulative percentages of children with a reported or substantiated maltreatment incidence increase substantially beyond the six months following a maltreatment incidence (i.e., 10.3% re-reporting after six months, 16.4% after twelve months, 20.6% after eighteen months). TIPS data for the NPP participants showed a similar pattern; thus, we chose the two-year time frame, as well, as this was the longest-term follow-up the data would allow.

2.5. Independent variables

2.5.1. Program dosage

Program dosage, our key variable of interest, was measured using the NPP data as a continuous variable of the total number of home and group sessions participated in by each caregiver.

2.5.2. Sociodemographic and caregiver characteristics

We controlled for sociodemographic characteristics of families associated with child maltreatment, including number of children, caregiver education, age, marital status, income, ethnicity, and gender (Hussey, Chang, & Kotch, 2006; Kotch et al., 1995; Miller & Cross, 2006; Wu et al., 2004). Demographic information was collected from participants before and after their participation in the NPP on the AAPI-2. In cases where pre- or post-test AAPI-2 demographic information was missing, we used information from the TIPS data. Dichotomous variables were constructed for gender (1 = female, 0 = male), and ethnicity (1 = non-white, 0 = white). We used the non-white categories to represent ethnicity since most of the participants (96%) were either African American or white with a small percentage of Hispanics, Native Americans, Pacific Islanders, and caregivers of other ethnicities, all of which were included in the non-white category. Caregiver age was coded continuously in years and marital status was measured by two dichotomous variables representing married (1 = married, 0 = not married) or cohabitating (1 = cohabitating, 0 = not cohabitating) with non-married and non-cohabitating (i.e., single) as the omitted category in the models. Number of children was coded as a continuous variable based on the number of children that caregivers reported. We constructed a dichotomous variable for whether or not the caregiver completed the 12th grade (1 = yes, 0 = no) based on their response to the highest grade level completed in school. (We do not have information on whether or not a caregiver received a GED or high school diploma – only years of schooling completed). This dichotomous variable was included in our model because in our exploratory analyses we did not find a linear relationship with years of education and our outcomes.

Rather, we found a significant association ($p < .05$) between some of our outcomes and 12th grade completion. Annual household income was measured as a continuous variable in thousands of dollars. This variable was constructed from income categories and converted to a continuous variable by using the mid point of the categories.

We also controlled for whether or not the caregiver experienced abuse within and/or outside their family as a child, given findings about the intergenerational transmission of maltreatment (Dorsey, Mustillo, Farmer, & Elbogen, 2008). These dichotomous variables were constructed from the AAPI-2 and measured whether or not the participant experienced any type of abuse outside of the family (1 = yes, 0 = no) and/or within the family (1 = yes, 0 = no).

We controlled for risk of maltreatment by including two measures. First, we used the AAPI-2 pre-test raw subscale scores, which measured parenting attitudes along the five dimensions outlined above in the description of the AAPI-2. These raw subscales are continuous measures, with higher scores indicating less risk. Second, we used the number of substantiated maltreatment investigations (from the TIPS data) for each caregiver from January 1, 1980 until the caregiver started the NPP. Since most of the prior investigations were due to neglect, we did not control for type of prior investigations. Finally, we included a set of dichotomous variables representing each resource center (with one center as the omitted category) to control for unmeasured differences between the centers.

2.6. Analysis

We used logistic regression to estimate the association between NPP attendance and the four outcome variables of interest – the presence of a reported or substantiated maltreatment incidence at six months or two years after participating in the program. While we explored different types of estimation approaches (count models) using different constructions of our dependent variables, the majority of individuals with maltreatment allegations, whether reported or substantiated, during the two years following a NPP closure date, had only one such allegation. Of those individuals with multiple allegations of maltreatment, reported or substantiated, within two years of the program end, the vast majority had only two allegations. Thus, logistic regression with a binary outcome was the most feasible approach. Finally, we used the robust clustering command in our logistic regression models in STATA 10 to adjust the standard errors for the lack of independence between observations due to the clustering of participants within centers (StataCorp, 2007).

2.7. Missing data

Across all variables in the model, 23% of the cases had some missing or incomplete information, primarily due to missing or incomplete information on the AAPI-2. To address the missing data, we used a multiple imputation routine in STATA 10 with the bootstrap option for all variables (Royston, 2007). We performed 25 imputations to match the percentage of missing data as recommended by Bodner (2008).

3. Results

3.1. Descriptive statistics

Table 1 presents descriptive statistics on the caregivers in the study population. In this table, we report descriptive statistics before and after multiple imputation to illustrate the similarity of the estimates and the extent of missing data for each variable. However, we focus on the imputed means in the text.

In terms of demographic characteristics, caregivers in our study population were 41.8% non-white and 74.2% female. Forty-three percent completed 12th grade and their average age was 27.4 years. Their average annual household income was \$13,900. Just over one-

Table 1
Study population characteristics of the Nurturing Parenting Program participants.

Characteristics	N	M (SD) or % before imputation	Range of values	M (SE) or % after imputation (N = 528)
Independent variables				
Program dosage	522	17.55 (9.69)	1–32	17.55 (.42)
Gender (%)	526			
Male		25.67		25.77
Female		74.33		74.23
Race/ethnicity (%)	526			
White		58.17		58.24
Non-white		41.83		41.76
Age at program start	526	27.38 (7.35)	12–60	27.38 (.32)
Income (in thousands)	452	13.92 (11.86)	7.5–60	13.88 (.57)
Educational level (%)	466			
Less than high school		55.80		57.45
High school or greater		44.20		42.55
Marital status (%)	501			
Single		63.48		63.37
Married		29.74		29.68
Cohabiting		6.78		6.95
Number of children	468	2.57 (1.50)	0–10	2.57 (.07)
Adult-Adolescent Parenting Inventory-2	442			
Inappropriate expectations		19.68 (4.58)	8–33	19.66 (.24)
Lack of empathy		36.92 (6.37)	19–50	36.96 (.31)
Corporal punishment		39.57 (7.69)	18–55	39.58 (.39)
Role reversal		23.39 (5.44)	9–35	23.22 (.27)
Power and independence		19.65 (3.00)	10–25	19.66 (.15)
Abuse as a child within family (%)	467	29.98		30.77
Abuse as a child outside of family (%)	467	18.63		18.63
Prior substantiated maltreatment investigations	528	1.20 (.99)	0–6	1.20 (.04)
Resource centers (% of sample)	528			
Provider 1		17.99		17.99
Provider 2		7.58		7.58
Provider 3		12.50		12.50
Provider 4		7.20		7.20
Provider 5		4.17		4.17
Provider 6		7.39		7.39
Provider 7		13.83		13.83
Provider 8		5.87		5.87
Provider 9		7.58		7.58
Provider 10		15.91		15.91
Dependent variables				
Reported incidence of maltreatment within six months of the NPP closure (%)	528	13.83		13.83
Substantiated incidence of maltreatment within six months of the NPP closure (%)	528	6.82		6.82
Reported incidence of maltreatment within two years of the NPP closure (%)	528	33.71		33.71
Substantiated incidence of maltreatment within two years of the NPP closure (%)	528	16.86		16.86

quarter were married (29.7%) and 7.0% were cohabitating. Caregivers had an average of 2.6 children.

Characteristics that may be associated with risk of maltreatment include history of abuse, prior maltreatment reports, and scores on the AAPI-2 subscales. Many of the caregivers were victims of abuse themselves as children. Almost one-third of the caregivers (30.8%) reported having experienced abuse as a child by a family member, and just under one-fifth (18.6%) from a perpetrator outside their family. Prior to being referred to the NPP, caregivers had an average of 1.2 substantiated reports of maltreatment. The means for caregivers' raw scores on the AAPI-2 sub-scales ranged from 19.7 (Inappropriate Parental Expectations of Children and Oppressing Children's Power and Independence) to 39.6 (Strong Belief in Corporal Punishment). (Higher scores indicate less risk.) Standardized to a population of abusive and non-abusive parents maintained by the instrument developer (Bavolek & Keene, 2001), the average population risk on each of the AAPI-2 sub-scales was considered high, except for the Strong Belief in Corporal Punishment, which was in the moderate risk category, but very close to the high risk cut-off. We used the raw scores in our analysis rather than the standardized risk scores because of their greater variation.

The caregivers participated in an average of 17.5 NPP group and home sessions across each of the ten resource centers. This average number of sessions is quite high given the sixteen-week program. It demonstrates the effectiveness of the supplemental home visits to maximize participation in the program and/or meet other family needs.

In terms of our dependent variable, at six months after participation in the NPP, 6.8% of the population had a substantiated and 13.8% had a reported maltreatment incidence after participating in the NPP. Two years following participation, 16.9% of the population had a substantiated maltreatment incidence compared to just over a third (33.7%) with a reported incidence.

3.2. Multivariate results

We used logistic regression to estimate the association between participation in the NPP and subsequent reported and substantiated maltreatment incidences at two different time points. This approach resulted in estimating four separate models as presented in Table 2. The coefficients for each resource center, with one as the omitted category, are not included in Table 2, but are available upon request. Many of the resource center coefficients in the models were

significant, suggesting unmeasured differences between the centers contribute to differences in subsequent maltreatment rates. We found differences by center in attendance, which ranged from 12 to 21 sessions, on average, for participants at each center. Yet, because we included attendance in our models as our key variable of interest, this alone does not explain differences in outcomes by center. Other unmeasured factors (i.e., model fidelity, staff characteristics) associated with both attendance and the outcomes could explain this observed relationship. First, we report results for the two outcomes (all reports and substantiated reports) at six months following NPP. Next, we report the two-year results.

3.3. Reported and substantiated maltreatment six months after the NPP

The first two models examined whether the degree of program attendance was associated with the likelihood of *all reported* and *substantiated* maltreatment incidences within a six-month time frame after participation in the program. The number of NPP sessions attended was significantly and negatively associated with the likelihood of all reported maltreatment incidences six months after participation in the program, but not substantiated incidences, though this relationship approaches significance in the same direction. More specifically, using post-estimation commands for categorical models developed by Long and Freese (2006) in STATA 10, we found that for each additional NPP session attended by a parent, the percent change in the odds of a reported incidence of maltreatment was approximately 3.8% lower, controlling for all other variables. Stated differently, for an otherwise average caregiver who completed three NPP sessions, the probability of having a subsequent maltreatment report was .177 compared to .108 for a caregiver who completed 18 sessions. These probabilities were calculated from the logistic regression models holding all other variables at their mean. These predicted probabilities are displayed in Fig. 1 along with the two-year results.

In the six-month models, only a few of our demographic and other control variables had a significant association with reported subsequent




Fig. 1. Likelihood of child maltreatment at two time periods after NPP participation.

maltreatment. Female caregivers and caregivers with more children were both significantly more likely than male caregivers or those with fewer children to have a reported incidence of subsequent maltreatment. Caregivers who graduated high school were significantly less likely than those who did not to have a subsequent maltreatment report during the six-month time period following the program.

In terms of substantiated reports, only caregiver's childhood experience of abuse by a family member had a significant association with the likelihood of a substantiated maltreatment in this time frame. More specifically, those caregivers who experienced abuse by a person within their family were significantly less likely to have a substantiated incidence of subsequent maltreatment within six months. While this finding may seem counterintuitive due to possible intergenerational transmission of maltreatment (Dorsey et al., 2008), the program may be particularly effective in changing the behavioral patterns of parents with this particular risk factor, given the premise of the program to target and change learned behaviors.

Table 2

Logistic regression models of reported and substantiated child maltreatment at two time periods after NPP completion (N = 528).

	Reported incidence within six months		Substantiated incidence within six months ^a		Reported incidence within two years		Substantiated incidence within two years	
	OR	SE	OR	SE	OR	SE	OR	SE
Program dosage	.96**	.01	.96	.02	.98	.01	.97**	.01
Female	1.90*	.49	1.54	.50	1.55	.39	1.28	.46
Nonwhite	.48	.19	.62	.16	.71	.20	.98	.48
Age	1.01	.02	1.01	.02	1.01	.01	1.00	.02
Income	1.01	.01	1.01	.02	1.01	.01	1.01	.01
High school graduate	.61*	.14	.89	.32	.69*	.11	.84	.20
Marital status ^b								
Married	.86	.27	.91	.43	1.08	.24	1.19	.46
Cohabiting	.59	.29	.68	.48	1.44	.80	1.74	1.03
Number of children	1.19*	.09	1.16	.15	1.27**	.10	1.19	.14
Adult-Adolescent Parenting Inventory-2								
Inappropriate expectations	.99	.04	.97	.07	.99	.03	1.01	.07
Lack of empathy	.98	.03	1.00	.04	.98	.03	.95	.03
Corporal punishment	1.03	.02	1.01	.04	1.02	.01	1.03	.02
Role reversal	.96	.03	.95	.05	.98	.02	.97	.03
Power and independence	.99	.04	.98	.09	.99	.04	1.06	.07
Abuse as a child (within family)	.67	.16	.32*	.18	.83	.28	.90	.35
Abuse as a child (outside of family)	1.33	.50	2.02	1.15	1.36	.33	1.25	.43
Prior substantiated maltreatment investigations	1.04	.18	1.13	.16	1.07	.13	1.19	.17

Note. The coefficients for each resource center were included in the models, but are not displayed in this table. They are available from the authors upon request.

^a In the logistic regression model with "substantiated incidence within six months" as the outcome variable, caregivers from one resource center were dropped from the analysis because there was no variation in the outcome variable, reducing the population size to 488.

^b Single is the reference category.

* p < .05.

** p < .01.

3.4. Reported and substantiated maltreatment two years after the NPP

In tracking the caregivers' Child Protective Services involvement two years after the program ended, a different pattern emerged. In fact, the amount of NPP participation only had a significant, negative relationship with *substantiated* maltreatment. For each additional NPP session parents attended, the odds of a substantiated reported incidence of subsequent maltreatment within two years decreased by 3.3%. As an illustration of effect size, for caregivers attending only three sessions of the NPP, the probability of a subsequent incidence of substantiated maltreatment within two years was .195, compared to a probability of .127 for those who attended 18, holding all other variables at their mean. These results are also displayed in Fig. 1.

As in the six-month models, only a few demographic characteristics predicted the likelihood of a *reported* incidence of subsequent maltreatment two years after the program ended. As the number of children increased, the likelihood of a reported incidence within this time frame significantly increased. Additionally, caregivers who completed 12th grade compared to those who did not were significantly less likely to have a subsequent maltreatment report within two years. None of the control variables, other than the resource centers, had a significant relationship with *substantiated* maltreatment incidences two years after participating in the NPP.

3.5. Foster care placement

We conducted some exploratory analyses to investigate whether the caregiver's child was in foster care at the time of participation in the NPP and whether this was a possible explanation for the different associations between program dosage and reported and substantiated maltreatment for the two time periods. Without controlling for any other variables, caregivers who had a child in foster care were less likely to have a subsequent reported and substantiated maltreatment incidence at six months after NPP participation and less likely to have a reported maltreatment incidence at two years post-NPP; this association disappeared when we controlled for prior risk and other demographic covariates. In other words, the presence of a child in foster care at a single time point did not affect the independent relationship between program dosage and the outcomes. We did not include the foster care variable in our final multivariate models because it does not adequately capture whether or not there may be another child in the home for whom a maltreatment incidence could occur or the timing and length of the foster care placement.

3.6. Multivariate results summary

In summary, the NPP was significantly associated with subsequent incidences of maltreatment, but the findings are dependent on both the timeframe during which the incidence occurred and the type of incidence — any reported or only substantiated incidences. Six months following participation, the number of sessions attended reduced the likelihood of reported incidences of maltreatment, but not substantiated. In the short-term, the program may positively change caregiver behavior in a great enough magnitude to raise fewer concerns by others, thus contributing to fewer overall re-reports. However, in the two-year time frame following participation, caregiver attendance in more sessions of the NPP was associated with fewer substantiated incidences of maltreatment, suggesting the program had enduring effects on the maltreatment outcomes of, perhaps, the greatest concern.

4. Discussion

Research on repeat maltreatment of children whose families are already involved in the child welfare system documents a substantial incidence of re-entry into foster care (Terling, 1999). Parent education is a primary vehicle used by caseworkers to help parents who strive to

keep their children from entering out-of-home care or who are seeking reunification. Evidence on the effectiveness of specific parent education programs on maltreatment outcomes for a child welfare population lags behind.

The results of this study, which build upon an initial evaluation of the NPP in Louisiana (Hodnett et al., 2009), provide several important findings about program effectiveness as it relates to the level of program participation. First, this study documents the significant association between the extent of program participation and fewer subsequent reports of child maltreatment within six months of program completion for families who have been referred to child welfare services. This finding is particularly important because it is well documented that maltreatment recurrences and re-referrals to child welfare agencies are most likely to occur within six months following a report of maltreatment (DePanfilis & Zuravin, 1999; English, Marshall, Brummel, & Orme, 1999; Fluke et al., 2005). Secondly, this study demonstrates the longer-term association between program participation and substantiated child maltreatment incidences. Long-term reductions in substantiated maltreatment incidences are the ultimate goal of any parent education program implemented in child welfare. And, our findings suggest a linkage between the NPP and the prevention of subsequent episodes of maltreatment and foster care re-entry, by extension, among families already involved in child welfare. This relationship, especially for those who participate more fully, appears to persist over time. Families involved in the child welfare system are often hard to engage in services, yet attendance at services has been shown to be negatively associated with the recurrence of child maltreatment (DePanfilis & Zuravin, 2002). As reported in our results, caregivers attended an average of 17.6 parenting sessions over a 16-week period, which represents a significant level of engagement and points to the relevance of this program and its resonance with families who are involved in the child welfare system. In addition, the average attendance levels demonstrate that coupling group programs with home-visiting outreach, as was done in Louisiana with the NPP, can achieve high levels of program participation, and thus, as demonstrated by our study, contribute to greater overall effectiveness.

Although a direct comparison cannot be made, the current study is similar to the study by Chaffin et al. (2004), in which PCIT was found to reduce repeat maltreatment in a physically abusive population. Of particular interest, however, is the fact that neglectful parents, who make up the majority of child welfare families as in our study population, did not demonstrate improvement after participation in PCIT. Further as a randomized control trial, significant resources were available to ensure the intervention was delivered with high quality and model integrity, neither of which are routinely achieved in a typical child welfare setting. In Louisiana, program fidelity monitoring was minimal to non-existent during the study period, though they have since put formal program observations and technical assistance in place for quality assurance purposes. Thus, the real world application of this program makes our findings for the NPP particularly impressive. Each additional program dose of the NPP was associated with reductions in subsequent maltreatment reports at six months or substantiations at two years after participation in the NPP by 3.8 and 3.3%, respectively.

Our study produced some results that require additional explanation — namely the different pattern of findings for the two types of reports (all reports vs. substantiated reports) at two time periods. These different patterns justify the need to examine reports and substantiation separately in order to identify and understand differential effects of interventions. To review, our results suggest that program dosage was only significantly associated with *reported* maltreatment incidences within six months and *substantiated* incidences within two years after participation in the NPP. Those caregivers who participated more intensely in the program, may be less likely to be reported, due to positive changes in behavior and/or service providers' perception of

these caregivers' willingness to change as compared to those who were less engaged. But, for the subsequent substantiated reports, which allegedly should be more severe, we did not detect a significant association in the short-term. This difference could also be the result of less power to detect differences given the smaller number of substantiated reports in the shorter time frame. Thus, while we found no differences in the number of substantiated incidences of maltreatment, greater program participation during this six-month time period, which is of particular interest to federal policy makers and the time period when repeat instances of maltreatment are most likely to occur, contributed to fewer overall reports of maltreatment.

Two years following program participation, our results indicate that caregivers appear to have made enough change in parenting behaviors to not engage in abusive or neglectful practices that would result in agency-verified substantiation. However, we saw no significant association with overall reports of maltreatment in the two-year time frame. Outside of child welfare involvement, the often unstable lifestyle of child welfare-involved families may still correspond with outside reports to child protective services. Attitudes, perceptions, and biases against higher risk caregivers may not be so easily changed regardless of a caregiver's improvement. Consequently, these findings suggest that while reporters continue to make allegations of abuse or neglect, upon investigation, those who benefited from attending more sessions were less likely to be substantiated for child maltreatment. Or, the program may only have enduring effects for families who are otherwise at risk for more serious maltreatment. Future research could elucidate the differential impact of common child welfare interventions, including parent education, on the type and timing of reports.

4.1. Limitations

Our study has several limitations in addition to its strengths. Since this study relies on a population of families served, our results are certainly generalizable to the child welfare population referred to the NPP in Louisiana; however, it may not be representative of other states or the national population. Even with a large study population such as this, we were unable to estimate the effect of NPP participation on type (i.e., abuse or neglect) or severity of repeat maltreatment due to the relatively low occurrence of certain types of maltreatment (i.e., physical and sexual abuse) on an already low frequency outcome (subsequent maltreatment) and the absence of data on severity.

Ideally, we would have liked to have limited our sample or adjusted the time frames for each case to account for the fact that some caregivers may not have had the "opportunity to maltreat" when their child(ren) were placed temporarily in out-of-home care. However, we were unable to use the administrative data to identify these time periods or identify the presence of other children in the home, even if one child was placed in out-of-home care. Thus, we were unable to accurately assess the "opportunity to maltreat," which means that some caregivers may have had less opportunity for repeat maltreatment than others.

In addition, we did find significant unmeasured differences in the outcomes by resource center, which we are unable to explain, though we did control for in our models. Future research should include measures of the NPP fidelity, which Louisiana is now putting into place and which may vary by center. It would also be worthwhile to explore other staff, service quality, and site-specific information and their associations with the outcomes.

Finally, one of the major limitations was our inability to definitively make causal attributions to NPP participation because other unmeasured factors that could be associated with program attendance might also explain the observed relationship with repeat maltreatment. In particular, an exploratory analysis revealed very few differences between high and low participators for the available variables included

in our models. We did find that caregivers who attended fewer sessions (less than the median of seventeen) were significantly more likely to have more prior substantiated reports than caregivers who attended more than the median, though we control for this difference in our models. This suggests, however, that it is certainly possible that other unmeasured differences in caregivers or the context in which they parent could explain the observed relationship between attendance and subsequent maltreatment, such as participant motivation, degree of economic hardship (Conger & Elder, 1997) or neighborhood factors (Garbarino & Kostelny, 1992). This form of self-selection bias as it relates to greater attendance warrants future investigation and is one that a large-scale experimental design could address.

5. Conclusion

The stakes for families involved in the child welfare system are high — child safety in the immediate sense and permanency and well-being for children in the long term. This study provides justification for continued use of the Nurturing Parenting Program for Infants, Toddlers, and Preschoolers in Louisiana and supports the need for ongoing evaluation of both process and outcome measures. By continuing to study promising models such as the NPP, we are adding to the evidence base of effective programs for this population. As promising evidence accumulates, the justification for investing significant resources in a randomized control trial of this program increases. These studies are needed to definitively establish a program's effectiveness with this population, which is, ultimately, what children and families deserve — investments that are known to work.

Acknowledgements

The authors would like to acknowledge Peter Pecora and Stephen Bavolek for helpful feedback and Jeremy Freese and Karen Grace-Martin for statistical consultation.

The current manuscript builds upon on a report by Hodnett et al. (2009), which analyzed some of the statewide data presented in this paper. This paper differs from the report in that it uses more data extracted over a longer period of time to examine the relationship between program dosage and subsequent maltreatment and employs statistical techniques to address missing data.

References

- Bandura, A. (1977). *Social learning theory*. Englewood Cliffs, NJ: Prentice Hall.
- Bandura, A. (1986). *Social foundations of thought and action: A social cognitive theory*. Englewood Cliffs, NJ: Prentice Hall.
- Barth, R. P. (1991). An experimental evaluation of in-home child abuse prevention services. *Child Abuse & Neglect*, 15(4), 363–375.
- Barth, R. P., Landsverk, J., Chamberlain, P., Reid, J. B., Rolls, J. A., Hurlburt, M. S., et al. (2005). Parent-training programs in child welfare services: Planning for a more evidence-based approach to serving biological parents. *Research on Social Work Practice*, 15(5), 353–371.
- Bavolek, S. (2005). Research and validation report of the Nurturing Parenting programs. Retrieved from http://www.nurturingparenting.com/research_validation/a4_main_r&v_article.pdf
- Bavolek, S., & Dellinger-Bavolek, J. (1985). Increasing the nurturing parenting skills of families in head start: Validation of the Nurturing Parenting program for parents and children birth to five years. Retrieved from http://www.nurturingparenting.com/research_validation/validation_b-5_program.pdf
- Bavolek, S. J., & Keene, R. G. (2001). *Adult-Adolescent Parenting Inventory-2: Administration and development handbook*. Asheville, NC: Family Development Resources, Inc.
- Bavolek, S. J., Kline, D. F., McLaughlin, J. A., & Publicover, P. R. (1979). Primary prevention of child abuse: Identification of high-risk adolescents. *Child Abuse & Neglect*, 3(3), 1071–1080.
- Bodner, T. E. (2008). What improves with increased missing data imputations? *Structural Equation Modeling*, 15(4), 651–675.
- Britner, P. A., & Reppucci, N. D. (1997). Prevention of child maltreatment: Evaluation of a parent education program for teen mothers. *Journal of Child and Family Studies*, 6(2), 165–175.
- Bugental, D. B., Ellerson, P. C., Lin, E. K., Rainey, B., Kokotovic, A., & O'Hara, N. (2002). A cognitive approach to child abuse prevention. *Journal of Family Psychology*, 16(3), 243–258.

- California Evidence-Based Clearinghouse for Child Welfare (2010). Nurturing Parenting Programs – summary. Retrieved from <http://www.cebc4cw.org/program/3>
- Chaffin, M., & Friedrich, B. (2004). Evidence-based treatments in child abuse and neglect. *Children and Youth Services Review*, 26(11), 1097–1113.
- Chaffin, M., Silovsky, J. F., Funderburk, B., Valle, L. A., Brestan, E. V., Balachova, T., et al. (2004). Parent-child interaction therapy with physically abusive parents: Efficacy for reducing future abuse reports. *Journal of Consulting and Clinical Psychology*, 72(3), 500–510.
- Conger, K. J., & Elder, G. H., Jr. (1997). Family economic hardship and adolescent adjustment: Mediating and moderating processes. In G. J. Duncan, & J. Brooks-Gunn (Eds.), *Consequences of growing up poor* (pp. 228–310). New York: Russell Sage Foundation.
- Cowen, P. S. (2001). Effectiveness of a parent education intervention for at risk families. *Journal of the Society for Pediatric Nurses*, 6(2), 73–82.
- Data Measures, Data Composites, and National Standards To Be Used in the Child and Family Services Reviews, 71 Fed. Reg. 32,969 (June 7, 2006).
- DePanfilis, D., & Zuravin, S. J. (1999). Epidemiology of child maltreatment recurrences. *The Social Service Review*, 73(2), 218–239.
- DePanfilis, D., & Zuravin, S. (2002). The effect of services on the recurrence of child maltreatment. *Child Abuse & Neglect*, 26(2), 187–205.
- Devall, E. (2004). Positive parenting for high-risk families. *Journal of Family and Consumer Sciences*, 96(4), 22–28.
- Donald, T., & Jureidini, J. (2004). Parenting capacity. *Child Abuse Review*, 13, 5–17.
- Dorsey, S., Mustillo, S. A., Farmer, E. M. Z., & Elbogen, E. (2008). Caseworker assessments of risk for recurrent maltreatment: Association with case-specific risk factors and re-reports. *Child Abuse & Neglect*, 32(3), 377–391.
- Drake, B., Jonson-Reid, M., Way, I., & Chung, S. (2003). Substantiation and recidivism. *Child Maltreatment*, 8(4), 248–260.
- Duggan, A., Caldera, D., Rodriguez, K., Burrell, L., Rohde, C., & Crowne, S. S. (2007). Impact of a statewide home visiting program to prevent child abuse. *Child Abuse & Neglect*, 31(8), 801–827.
- Duggan, A., McFarlane, E., Fuddy, L., Burrell, L., Higman, S. M., Windham, A., et al. (2004). Randomized trial of a statewide home visiting program: Impact in preventing child abuse and neglect. *Child Abuse & Neglect*, 28(6), 597–622.
- Duggan, A. K., McFarlane, E. C., Windham, A. M., Rohde, C. A., Salkaver, D. S., Fuddy, L., et al. (1999). Evaluation of Hawaii's Healthy Start Program. *The Future of Children*, 9(1), 66–90.
- DuMont, K., Mitchell-Herzfeld, S., Greene, R., Lee, E., Lowenfels, A., Rodriguez, M., et al. (2008). Healthy Families New York (HFNY) randomized trial: Effects on early child abuse and neglect. *Child Abuse & Neglect*, 32(3), 295–315.
- English, D. J., Marshall, D. B., Brummel, S., & Orme, M. (1999). Characteristics of repeated referrals to child protective services in Washington state. *Child Maltreatment*, 4(4), 297–307.
- Erickson, M. F., & Egeland, B. (2011). Child neglect. In J. E. B. Myers (Ed.), *The APSAC handbook on child maltreatment* (pp. 103–124). (3rd ed.). Thousand Oaks, CA: Sage Publications, Inc.
- Eyberg, S. M., & Robinson, E. A. (1982). Parent-child interaction therapy: Effects on family functioning. *Journal of Clinical Child Psychology*, 11, 130–137.
- Fluke, J. D., Shusterman, G. R., Hollinshead, D., & Yuan, Y. T. (2005). *Reporting and recurrence of child maltreatment: Findings from NCANDS*. Washington, DC: U.S. Department of Health and Human Services, Office of the Assistant Secretary for Planning and Evaluation.
- Garbarino, J., & Kostelny, K. (1992). Child maltreatment as a community problem. *Child Abuse & Neglect*, 16(4), 455–464.
- Gershater-Molko, R. M., Lutzker, J. R., & Wesch, D. (2002). Using recidivism data to evaluate Project Safecare: Teaching bonding, safety, and health care skills to parents. *Child Maltreatment*, 7(3), 277–285.
- Gilbert, R., Widom, C. S., Browne, K., Fergusson, D., Webb, E., & Janson, S. (2009). Burden and consequences of child maltreatment in high-income countries. *Lancet*, 373(9657), 68–81.
- Haapasalo, J., & Pokela, E. (1999). Child-rearing and child abuse antecedents of criminality. *Aggression and Violent Behavior*, 4(1), 107–127.
- Henggeler, S. W., Schoenwald, S. K., Borduin, C. M., Rowland, M. D., & Cunningham, P. B. (1998). *Multisystemic treatment of antisocial behavior in children and adolescents*. New York: Guilford Press.
- Hodnett, R., Faulk, K., Dellinger, A., & Maher, E. (2009). *Evaluation of the statewide implementation of a parent education program in Louisiana's child welfare agency: The Nurturing Parenting Program for infants, toddlers, and pre-school children*. Seattle, WA: Casey Family Programs.
- Huebner, C. (2002). Evaluation of a clinic-based parent education program to reduce the risk of infant and toddler maltreatment. *Public Health Nursing*, 19(5), 377–389.
- Hurlburt, M., Barth, R., Leslie, L., & Landsverk, J. (2007). Building on strengths: Current status and opportunities for improvement of parent training for families in child welfare. In R. Haskins, F. Wulczyn, & M. B. Webb (Eds.), *Child protection: Using research to improve policy and practice* (pp. 81–106). Washington, DC: Brookings Institute Press.
- Hussey, J. M., Chang, J. J., & Kotch, J. B. (2006). Child maltreatment in the United States: Prevalence, risk factors, and adolescent health consequences. *Pediatrics*, 118(3), 933–942.
- Hussey, J. M., Marshall, J. M., English, D. J., Knight, E. D., Lau, A. S., Dubowitz, H., et al. (2005). Defining maltreatment according to substantiation: Distinction without a difference? *Child Abuse & Neglect*, 29(5), 479–492.
- Huxley, P., & Warner, R. (1993). Primary prevention of parenting dysfunction in high-risk cases. *The American Journal of Orthopsychiatry*, 63(4), 582–588.
- Johnson, M. A., Stone, S., Lou, C., Ling, J., Claassen, J., & Austin, M. J. (2008). Assessing parent education programs for families involved with child welfare services: Evidence and implications. *Journal of Evidence-Based Social Work*, 5(1/2), 191–236.
- Kaminski, J. W., Valle, L. A., Filene, J. H., & Boyle, C. L. (2008). A meta-analytic review of components associated with parent training program effectiveness. *Journal of Abnormal Child Psychology*, 36(4), 567–589.
- Kochanska, G., & Aksan, N. (1995). Mother-child mutually positive affect: The quality of child compliance to requests, prohibitions, and maternal control as correlates of early internalization. *Child Development*, 66, 236–254.
- Kotch, J. B., Browne, D. C., Ringwalt, C. L., Stewart, P. W., Ruina, E., Holt, K., et al. (1995). Risk of child abuse or neglect in a cohort of low-income children. *Child Abuse & Neglect*, 19(9), 1115–1130.
- Laible, D. (2004). Mother-child discourse in two contexts: Links with child temperament, attachment security, and socioemotional competence. *Developmental Psychology*, 40(6), 979–992.
- Lee, S., Aos, S., & Miller, M. (2008). *Evidence-based programs to prevent children from entering and remaining in the child welfare system: Benefits and costs for Washington (Document No. 08-07-3901)*. Olympia, WA: Washington State Institute for Public Policy.
- Letarte, M. -J., Normandeau, S., & Allard, J. (2010). Effectiveness of a parent training program "Incredible Years" in a child protection service. *Child Abuse & Neglect*, 34(4), 253–261.
- Littell, J. H. (1997). Effects of the duration, intensity, and breadth of family preservation services: A new analysis of data from the Illinois Family First experiment. *Children and Youth Services Review*, 19(1/2), 17–39.
- Long, J. S., & Freese, J. (2006). *Regression models for categorical dependent variables using Stata* (2nd ed.). College Station, TX: Stata Press.
- Lundahl, B. W., Nimer, J., & Parsons, B. (2006). Preventing child abuse: A meta-analysis of parent training programs. *Research on Social Work Practice*, 16(3), 251–262.
- MacMillan, H. L., Thomas, B. H., Jamieson, E., Walsh, C. A., Boyle, M. H., Shannon, H. S., et al. (2005). Effectiveness of home visitation by public-health nurses in prevention of the recurrence of child physical abuse and neglect: A randomised controlled trial. *Lancet*, 365(9473), 1786–1793.
- Marcenko, M. O., Spence, M., & Samost, L. (1996). Outcomes of a home visitation trial for pregnant and postpartum women at-risk for child placement. *Children and Youth Services Review*, 18(3), 243–259.
- Marcynyszyn, L. A., Maher, E. J., Corwin, T. W. (2011). Getting with the (evidence-based) program: An evaluation of the Incredible Years Parent Training Program in child welfare. *Children and Youth Services Review*, 33(5), 747–757.
- Miller, A. B., & Cross, T. (2006). Ethnicity in child maltreatment research: A replication of Behl et al.'s content analysis. *Child Maltreatment*, 11(1), 16–26.
- Olds, D. L., Eckenrode, J., Henderson, C. R., Kitzman, H., Powers, J., Cole, R., et al. (1997). Long-term effects of home visitation on maternal life course and child abuse and neglect: Fifteen-year follow-up of a randomized trial. *Journal of the American Medical Association*, 278(8), 637–643.
- Patterson, G. R., Reid, J. B., & Eddy, J. M. (2002). A brief history of the Oregon model. In J. B. Reid, G. R. Patterson, & J. J. Snyder (Eds.), *Antisocial behavior in children and adolescents: A developmental analysis and model for intervention* (pp. 3–20). Washington, DC: American Psychological Association.
- Prinz, R. J., Sanders, M. R., Shapiro, C. J., Whitaker, D. J., & Lutzker, J. R. (2009). Population-based prevention of child maltreatment: The U.S. Triple P System population trial. *Prevention Science*, 10(1), 1–12.
- Reynolds, A. J., Mathieson, L. C., & Topitzes, J. W. (2009). Do early childhood interventions prevent child maltreatment?: A review of research. *Child Maltreatment*, 14(2), 182–206.
- Royston, P. (2007). Multiple imputation of missing values: Further update of ice, with an emphasis on interval censoring. *STATA Journal*, 7, 445–464.
- StataCorp (2007). *Stata Statistical Software: Release 10*. College Station, TX: StataCorp LP.
- Stern, S. B., Alaggia, R., Watson, K., & Morton, T. R. (2008). Implementing an evidence-based parenting program with adherence in the real world of community practice. *Research on Social Work Practice*, 18(6), 543–554.
- Substance Abuse and Mental Health Services Administration's National Registry of Evidence-Based Programs and Practices (2010). Nurturing Parenting Programs. Retrieved from <http://www.nrepp.samhsa.gov/ViewIntervention.aspx?id=171>
- Terling, T. (1999). The efficacy of family reunification practices: Reentry rates and correlates of reentry for abused and neglected children reunited with their families. *Child Abuse & Neglect*, 23(12), 1359–1370.
- Thomlison, B. (2005). Using evidence-based knowledge to improve policies and practices in child welfare: Current thinking and continuing challenges. *Research on Social Work Practice*, 15(5), 321–322.
- U.S. Department of Health and Human Services, Administration for Children and Families, Administration on Children, Youth and Families, Children's Bureau (2010). *Child maltreatment 2009*. Washington, DC: U.S. Government Printing Office.
- Wagner, M. M., & Clayton, S. L. (1999). The Parents as Teachers Program: Results from two demonstrations. *The Future of Children*, 9(1), 91–115.
- Waldfogel, J. (2009). Prevention and the child protection system. *The Future of Children*, 19(2), 195–210.
- Webster-Stratton, C. (2007). *The Incredible Years parent training manual: BASIC Program* (3rd ed.). Seattle, WA: Incredible Years.
- Widom, C. S., & Maxfield, M. G. (2001). *An update on the "cycle of violence."* Research in brief (NCJ Publication No. 184894). Washington, D.C.: U.S. Dept. of Justice, Office of Justice Programs, National Institute of Justice.
- Wu, S. S., Ma, C. -X., Carter, R. L., Ariet, M., Feaver, E. A., Resnick, M. B., et al. (2004). Risk factors for infant maltreatment: A population-based study. *Child Abuse & Neglect*, 28(12), 1253–1264.
- Wulczyn, F., Hislop, K., & Jones Harden, B. (2002). The placement of infants in foster care. *Infant Mental Health Journal*, 23(5), 454–475.